

Supplementary Online Content

Secemsky EA, Yeh RW, Kereiakes DJ, et al; DAPT Study Investigators. Mortality following cardiovascular and bleeding events occurring beyond 1 year after coronary stenting: a secondary analysis of the Dual Antiplatelet Therapy (DAPT) Study. *JAMA Cardiol*. Published online March 15, 2017. doi:10.1001/jamacardio.2017.0063

eTable 1. Risk of Mortality After Ischemic and Bleeding Events During the 21-Month Post-Randomization Period, Stratified by <30 or ≥30 Days From Time of Event

eTable 2. Frequency of Bleeding Events and Death by BARC Classification

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Risk of Mortality After Ischemic and Bleeding Events During the 21-Month Postrandomization Period, Stratified by ≤ 30 or >30 Days From Time of Event

Endpoints	Adjusted HR for Mortality,* ≤ 30 days (95% CI)	Adjusted HR for Mortality,* >30 days (95% CI)
Ischemic Events		
MI not related to ST	33.0 (17.8-61.3)	5.8 (3.6-9.4)
Definite or probable ST	135.8 (75.1-245.3)	1.0 (0.1-6.9)
Ischemic stroke	95.7 (48.1-190.4)	3.4 (1.1-10.7)
Any MI, definite or probable ST, or ischemic stroke	72.0 (48.8-106.4)	4.1 (2.5-6.5)
Bleeding Events		
GUSTO moderate bleed	40.5 (18.8-87.2)	4.4 (2.1-9.0)
GUSTO severe bleed	230.8 (136.6-389.8)	10.1 (4.9-20.9)
GUSTO moderate/severe bleed	103.4 (66.9-159.8)	6.3 (3.7-10.6)
BARC 2, 3, or 5 bleed	53.4 (34.7-82.0)	3.8 (2.4-6.0)
BARC 2 or 3 bleed	20.8 (11.2-38.5)	3.7 (2.3-5.9)
BARC 3 or 5 bleed	92.5 (59.5-143.6)	5.6 (3.3-9.6)
BARC 2 bleed	10.4 (3.3-32.6)	2.7 (1.4-5.3)
BARC 3 bleed	31.7 (16.1-62.6)	5.5 (3.2-9.5)
*Adjusted models were stratified by propensity score quintile. Propensity scores were derived from all variables included in Tables 1 and 2 in the article. Abbreviations: BARC, Bleeding Academic Research Consortium; GUSTO, Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Arteries; HR, hazard ratio; MI, myocardial infarction; ST, stent thrombosis.		

eTable 2. Frequency of Bleeding Events and Death by BARC Classification

Bleeding Endpoint, No. (%)	Events after Randomization *	Rates of Death After an Event	Cumulative Incidence of Death After an Event
BARC 2, 3, or 5	481 (4.1)	46 (9.56)	0.42
BARC 3 or 5	244 (2.1)	39 (15.98)	0.36
BARC 2	262 (2.3)	12 (4.58)	0.11
BARC 3	229 (1.97)	24 (10.48)	0.22
BARC 5	15 (0.13)	15 (100.00)	0.14
<p>*The number of each event represents the number of patients with the event. Multiple types of events could be accrued by the same patient. For patients with >1 of the same type of event, only the earliest event contributed to the analysis. Abbreviation: BARC, Bleeding Academic Research Consortium</p>			