

Supplementary Online Content

Idorn LW, Datta P, Heydenreich J, Philipsen PA, Wulf HC. A 3-year follow-up of sun behavior in patient with cutaneous malignant melanoma. Published online October 2, 2013. *JAMA Dermatol*. doi: 10.1001/jamadermatol.2013.5098.

eTable. Sun Diary Data With Information About Various Exposure Situations

This supplementary material has been provided by the authors to give readers additional information about their work.

e-Table. Sun Diary Data With Information About Various Exposure Situations^a

Variable	No. of Days, Mean (SD)			
	Group 1 First Summer After Diagnosis (2010)	Group 1 Second Summer After Diagnosis (2011)	Group 2 First Summer After Diagnosis (2009)	Group 2 Third Summer After Diagnosis (2011)
With dosimeter measurements and diary information				
Patients	80 ± 16	74 ± 3	69 ± 12	70 ± 9
Controls	74 ± 12	72 ± 5	78 ± 4	74 ± 6
At work				
Patients	39 ± 16	34 ± 13	25 ± 19	27 ± 16
Controls	28 ± 15	33 ± 12	23 ± 27	26 ± 17
Off work				
Patients	27 ± 9	30 ± 19	42 ± 28	35 ± 21
Controls	32 ± 14	28 ± 12	48 ± 26	37 ± 20
On holiday ^b				
Patients	14 ± 12	10 ± 8	3 ± 4	9 ± 8
Controls	15 ± 12	12 ± 8	7 ± 7	10 ± 8
Abroad				
Patients	11 ± 10	10 ± 8	1 ± 3	4 ± 7
Controls	2 ± 4	2 ± 4	5 ± 4	4 ± 4
Sunburned (red from the sun)				
Patients	2 ± 2	1 ± 2	2 ± 2	1 ± 1
Controls	2 ± 2	2 ± 2	1 ± 2	1 ± 2
Using sunscreen				
Patients	29 ± 28	18 ± 21	28 ± 23	28 ± 25
Controls	14 ± 21	14 ± 23	15 ± 24	19 ± 27
With body exposure ^c				
Patients	10 ± 12	6 ± 10	10 ± 10	11 ± 11
Controls	15 ± 12	10 ± 12	8 ± 6	9 ± 5
With body exposure without using sunscreen				
Patients	1 ± 1	1 ± 3	2 ± 3	3 ± 5

Controls	9 ± 8	6 ± 9	4 ± 5	6 ± 5
----------	-------	-------	-------	-------

Abbreviation: SED, standard erythema dose.

^aGroups A and B each comprised 10 patients and 10 matched controls.

^bAway from home during a holiday.

^cSunbathing, exposing the shoulders or upper body outdoors, or using a solarium.