

Supplementary Online Content

Nosrati A, Berliner JG, Goel S, et al. Outcomes of melanoma in situ treated with Mohs micrographic surgery compared with wide local excision. *JAMA Dermatol*. Published online February 22, 2017. doi:10.1001/jamadermatol.2016.6138

eTable. Procedures performed by year

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Procedures performed by year			
Year of diagnosis	Procedure	Frequency	Percent*
1978	WLE	1	0.15
1985	WLE	1	0.15
1986	MMS	1	0.15
1988	WLE	3	0.45
1989	WLE	1	0.15
1990	MMS	1	0.15
1990	WLE	3	0.45
1991	WLE	4	0.60
1992	WLE	6	0.91
1993	WLE	5	0.76
1994	WLE	9	1.36
1995	WLE	4	0.60
1996	MMS	1	0.15
1996	WLE	7	1.06
1997	WLE	14	2.11
1998	MMS	1	0.15
1998	WLE	16	2.42
1999	WLE	17	2.57
2000	MMS	4	0.60
2000	WLE	36	5.44
2001	MMS	7	1.06
2001	WLE	19	2.87
2002	MMS	14	2.11
2002	WLE	27	4.08
2003	MMS	15	2.27
2003	WLE	53	8.01
2004	MMS	28	4.23
2004	WLE	44	6.65
2005	MMS	29	4.38
2005	WLE	13	1.96
2006	MMS	12	1.81
2006	WLE	4	0.60
2007	MMS	18	2.72
2007	WLE	15	2.27
2008	MMS	31	4.68
2008	WLE	16	2.42
2009	MMS	29	4.38
2009	WLE	16	2.42
2010	MMS	22	3.32

2010	WLE	12	1.81
------	-----	----	------

Year of diagnosis	Procedure	Frequency	Percent*
2011	MMS	23	3.47
2011	WLE	8	1.21
2012	MMS	32	4.83
2012	WLE	23	3.47
2013	MMS	8	1.21
2013	WLE	8	1.21

Mohs micrographic surgery; WLE, wide local excision

*Percent refers to percent of all procedures