Supplementary Online Content

Stolmeier DA, Stratman HB, McIntee TJ, Stratman EJ. Utility of laboratory test result monitoring in patients taking oral terbinafine or griseofulvin for dermatophyte infections. *JAMA Dermatol.* Published online October 17, 2018. doi:10.1001/jamadermatol.2018.3578

eTable. Common Terminology Criteria for Adverse Events

This supplementary material has been provided by the authors to give readers additional information about their work.

© 2018 American Medical Association. All rights reserved.

eTable. Common Terminology Criteria for Adverse Events

Adverse Event	Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
Alanine aminotransferase (ALT) increased	>ULN – 3.0 x ULN Adults: >ULN – 120 Children: >ULN – 66	>3.0 – 5.0 x ULN Adults: 121 – 200 Children: 67–110	>5.0 – 20.0 x ULN Adults: 201– 800 Children: 111–440	>20.0 x ULN Adults: >800 Children:>440	-
Aspartate aminotransferase (AST) increased	>ULN – 3.0 x ULN Adults: >ULN – 120 Children: >ULN – 96	>3.0 – 5.0 x ULN Adults: 121 – 200 Children: 97 – 160	>5.0 – 20.0 x ULN Adults: 201– 800 Children: 161–640	>20.0 x ULN Adults: >800 Children: >640	-
Anemia	Hemoglobin (Hgb) < LLN – 10.0 g/dL	Hgb <10.0 – 8.0 g/dL	Hgb <8.0 g/dL	Life-threatening consequences; urgent intervention indicated	Death
Neutrophil count decreased	<LLN – 1.5 x 10e9 /L	<1.5 – 1.0 x 10e9 /L	<1.0 – 0.5 x 10e9 /L	<0.5 x 10e9 /L	-
Lymphocyte count decreased	<LLN – 0.8 x 10e9 /L	<0.8 – 0.5 x 10e9 /L	<0.5 -0.2 x 10e9 /L	<0.2 x 10e9 /L	-

© 2018 American Medical Association. All rights reserved.