

Supplementary Online Content


Lortet-Tieulent J, Goding Sauer A, Siegel RL, et al. State-level cancer mortality attributable to cigarette smoking in the United States. *JAMA Intern Med*. Published online October 24, 2016. doi:10.1001/jamainternmed.2016.6530

eFigure. State proportion of cancer deaths attributable to cigarette smoking, in non-Hispanic whites and in all races and ethnicities, in men, in 2014

eTable. Number and proportion of cancer deaths attributable by US region to smoking using smoking prevalence from the Behavioral Risk Factor Surveillance System (BRFSS) and the National Health Interview Survey (NHIS)

This supplementary material has been provided by the authors to give readers additional information about their work.

eFigure. State proportion of cancer deaths attributable to cigarette smoking, in non-Hispanic whites and in all races and ethnicities, in men, in 2014



eTable. Number and proportion of cancer deaths attributable by US region to smoking using smoking prevalence from the Behavioral Risk Factor Surveillance System (BRFSS) and the National Health Interview Survey (NHIS)

	Cancer deaths attributable to smoking and 95% confidence interval		Proportion of cancer deaths attributable to smoking (%) and 95% confidence interval	
	BRFSS	NHIS	BRFSS	NHIS
Northeast	29698	32312	27.3 (25.9;28.6)	28.2 (26.4;29.8)
Midwest	40164	40235	29.3 (27.9;30.6)	29.5 (27.9;30.9)
South	66069	65375	29.8 (28.4;31.1)	30.3 (29.0;31.5)
West	32037	30543	26.4 (25.0;27.8)	27.3 (25.7;28.7)
US	167968	168465	28.7 (27.8;29.2)	28.8 (28.2;30.2)