

Supplementary Online Content



Appendix 14: Determination of Brain Death/Death by Neurologic Criteria (BD/DNC)


Process Overview



Based on recommendations developed by the World Brain Death Project (see main paper in JAMA)


To be used with the Determination of BD/DNC Checklist for Use in Adults and Children ≥ 36 Weeks of Age (see Supplement 15)

Section 1.a. Assess for Prerequisites and Confounders*




^{*}Refer to Supplement 15: Prerequisites for Clinical Examination, from the Checklist for Determination of Brain Death/Death by Neurologic Criteria

Section 1.b. Assess for Prerequisites and Confounders in Persons Treated with Therapeutic Hypothermia*



^{*}Refer to Supplement 15: Prerequisites for Clinical Examination, from the Checklist for Determination of Brain Death/Death by Neurologic Criteria

Section 2. Perform Clinical Exam*




Ancillary testing is required when:


- There are confounding conditions that cannot be resolved,
- The clinical exam has been completed to the fullest extent possible and there is an inability to complete all tests,
- Uncertainty exists regarding interpretation of spinal-mediated motor responses,
- The whole brain death formulation is being followed, and there is isolated brainstem pathology,
- Dictated by law or regional guidelines.

^{*}Refer to Supplement 15: Prerequisites for Clinical Examination, from the Checklist for Determination of Brain Death/Death by Neurologic Criteria

Section 3. Perform Apnea Testing*



Section 4. Perform Ancillary Testing, as Required*



^{*}Refer to Supplement 15: Acceptable Ancillary Testing Modallities, from the Checklist for Determination of Brain Death/Death by Neurologic Criteria