

Supplementary Online Content

Jernberg T, Johanson P, Held C, Svennblad B, Lindbäck J, Wallentin L; SWEDHEART/RIKS-HIA. Association between adoption of evidence-based treatment and survival for patients with ST-elevation myocardial infarction. *JAMA*. 2011;305(16):1677-1684.

eTable 1. Observed (Crude) and Standardized Mortality (%) When Only Hospitals Participating During the Whole Study Period Were Included in the Analysis

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Observed (Crude) and Standardized Mortality (%) When Only Hospitals Participating During the Whole Study Period Were Included in the Analysis

	In-hospital		30-day		1-year	
	Observed	Standardized	Observed	Standardized	Observed	Standardized
1996	11.5	10.6	14.1	12.9	20.0	18.8
1997	11.0	10.6	13.7	13.2	19.5	19.6
1998	10.4	9.5	13.4	12.2	19.2	18.6
1999	10.3	9.3	12.7	11.5	18.2	17.4
2000	9.4	8.2	11.9	10.3	17.5	16.3
2001	8.6	7.8	10.6	9.7	16.7	16.0
2002	8.5	7.7	11.2	10.4	16.1	15.6
2003	7.4	7.6	8.9	9.2	14.0	14.8
2004	7.0	6.9	9.2	9.1	13.3	13.9
2005	6.1	5.9	7.8	7.6	13.1	13.3
2006	5.7	5.7	7.1	7.2	11.2	11.4
2007	5.3	5.3	6.6	6.6	11.2	11.2

Standardized mortality is expected number of deaths that would have occurred if the distribution of baseline characteristics were the same for all years as in 2007.