

Supplementary Online Content

Li J, Cone JE, Kahn AR. Association between World Trade Center exposure and excess cancer risk, 2003-2008 *JAMA*. doi:10.1001/jama.2012.110980.

eTable. Definition of World Trade Center (WTC)-related exposures*

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Definition of World Trade Center (WTC)-related exposures*

WTC-related exposure level	Rescue and recovery exposure among rescue/recovery workers	WTC-related residential, office, school, or pass-through exposure among participants not involved in rescue/recovery
High	Workers who were in Manhattan, south of Chambers Street, between the time of the first plane impact and 1200 h on 9/11 (encompassing the collapse of WTC towers) and who worked on the WTC dust and debris pile on 9/11 or worked at the WTC site for more than 90 days, starting before Sept 18, 2001.	Participants who reported two or more injuries on 9/11; and on 9/11 (a) worked or was a passerby south of Chamber street in lower Manhattan, (b) resided in Lower Manhattan zip codes including Canal Street and south and did not evacuate their home, or (c) were present at school in Lower Manhattan zip codes including Canal Street and south.
Low	Workers who began work at the WTC site after Sept 17, 2001, did not work on the WTC debris pile, worked fewer than 30 days at the WTC site, and were not present south of Chambers Street between the first plane impact and 1200 h on 9/11.	Participants who reported no injuries on 9/11, and on 9/11 (a) worked or was a passerby below south of Chamber street, (b) resided in Lower Manhattan zip codes including Canal Street and south, and evacuated their home for any amount of time, or (c) were not present at school in Lower Manhattan zip codes including Canal Street and south.
Intermediate	All other rescue/recovery workers.	All other participants not involved in rescue/recovery.

*Adapted from previous study by HT Jordan (*Lancet* 2011;378:879-87)