

Supplementary Online Content

Warner DO, Berge K, Sun H, Harman A, Hanson A, Schroeder DR. Substance use disorder among anesthesiology residents, 1975-2009. *JAMA*. doi:10.1001/jama.2013.281954

eTable. Estimated Proportion of Residents Who Relapsed by Year Apparent

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Estimated Proportion of Residents Who Relapsed by Year Apparent*

Year apparent	Years after apparent	Estimate†	95% CI
1975-1984	1	10%	(0-27)
	2	10%	(0-27)
	5	20%	(0-41)
	10	30%	(0-53)
	15	50%	(7-73)
	20	60%	(15-81)
	25	60%	(15-81)
1985-1994	1	7%	(2-11)
	2	8%	(2-13)
	5	15%	(8-22)
	10	24%	(15-32)
	15	34%	(24-42)
	20	38%	(27-47)
	25	40%	(29-50)
1995-2004	1	11%	(5-17)
	2	17%	(9-24)
	5	29%	(20-37)
	10	32%	(22-41)
	15	39%	(25-50)
2005-2010	1	8%	(3-14)
	2	11%	(4-17)
	5	15%	(7-23)

* N=310 (91 events) included in analysis.

† Kaplan-Meier estimate for number who relapsed.