

Supplementary Online Content

El Khamali R, Mouaci A, Valera S, et al; for the SISTRESSREA Study Group. Effects of a multimodal program including simulation on job strain among nurses working in intensive care units: a randomized clinical trial. *JAMA*. doi:10.1001/jama.2018.14284 Published online October 24, 2018.

eTable. Prevalence of high psychological demand, low decision latitude and low social support on inclusion and at 6 months (secondary outcomes)

This supplementary material has been provided by the authors to give readers additional information about their work.

© 2018 American Medical Association. All rights reserved.

eTable. Prevalence of high psychological demand, low decision latitude and low social support on inclusion and at 6 months (secondary outcomes)

	Inclusion			At 6 months		
	Intervention N = 101	Control N = 97	Difference between groups (95% CI)	Intervention N = 97	Control N = 85	Difference between groups (95% CI)
High psychological demand, N (%)	79 (78)	67 (69)	9% [95%CI, -3 to 21]	26 (27)*	61 (72)	-45% [95%CI, -56 to -31]
Low decision latitude, N (%)	74 (73)	63 (65)	8% [95%CI, -5 to 20]	15 (15)*	58 (68)	-53% [95%CI, -63 to -39]
Low social support, N (%)	65 (64)	61 (63)	1% [95%CI, -12 to 15]	49 (51)*	65 (76)	-26% [95%CI, -38 to -12]

* $P < 0.001$ vs. control by Pearson chi-square test