

Supplementary Online Content

Goldman-Mellor S, Olfson M, Lidon-Moyano C, Schoenbaum M. Association of suicide and other mortality with emergency department presentation. *JAMA Netw Open*. 2019;2(12):e1917571. doi:10.1001/jamanetworkopen.2019.17571

eTable. ICD-9-CM Codes Used to Define Method of Self-Harm Injury


eFigure. Kaplan-Meier Cumulative Probabilities of Suicide for 1 Year of Follow-Up During 2009 to 2012 Among California Emergency Department Patients Stratified by Patient Group and Sex

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. ICD-9-CM Codes Used to Define Method of Self-Harm Injury

ICD-9-CM codes used to define method of self-harm injury.	
Category	ICD-9-CM codes
Poisoning	E950.0-E952.x
Cut/pierce	E956.0-E956.x
Hanging/strangulation/suffocation	E953.0-E953.x
Jumping	E957.0-E957.x
Firearm	E955.0-E955.x
Other	E954.0-E954.x, E958.0-E958.x

eFigure. Kaplan-Meier Cumulative Probabilities of Suicide for 1 Year of Follow-Up During 2009 to 2012 Among California Emergency Department Patients Stratified by Patient Group and Sex



eFigure. Kaplan-Meier cumulative probabilities of suicide, by patient group and sex, for 365 days of follow-up during 2009-2012 among California emergency department patients. Log-rank tests comparing suicide rates among male vs. female patients were significant in all patient groups (all $p < 0.001$).