

Supplementary Online Content

Castrioto A, Lozano AM, Poon Y-Y, Lang AE, Fallis M, Moro E. Ten-year outcome of subthalamic stimulation in Parkinson disease: a blinded evaluation. *Arch Neurol*. Published online August 8, 2011. doi:10.1001/archneurol.2011.182.

eTable. Adverse events reported at 1-, 5- and 10-year follow-up.

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Adverse events reported at 1-, 5- and 10-year follow-up.

Adverse events	1 year No. (%)	5 years No. (%)	10 years No. (%)
Hypophonia*	11 (61)	11 (61)	11 (61)
Weight gain	12 (66.6)	12 (66.6)	8 (44.4)
Eyelid opening apraxia*	5 (27.8)	5 (27.8)	5 (27.8)
Cognitive decline	0	2 (11.1)	4 (22.2)
Infection	2 (11.1)	0	2 (11.1)
Depression†	4 (22.2)	3 (16.7)	3 (16.7)
Impulse control disorders†	NA	3 (16.7)	3 (16.7)
Visual hallucinations	0	0	4 (22.2)
Dysphagia	0	0	2 (11.1)

Abbreviation: NA, not available.

* For hypophonia and eyelid opening apraxia no change in symptoms between 1-year, 5-year and 10-year assessments.

† For depression and impulse control disorders there was no change in symptoms between the 5-year and 10-year assessments.