

Supplementary Online Content

Murphy JE, Wo JY, Ryan DP, et al. Total neoadjuvant therapy with FOLFIRINOX followed by individualized chemoradiotherapy for borderline resectable pancreatic adenocarcinoma. *JAMA Oncol*. Published online May 3, 2018.
doi:10.1001/jamaoncol.2018.0329

eTable. Pathologic Characteristics at Surgery of Resected Patients

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Pathologic Characteristics at Surgery of Resected Patients

| | All Eligible (N=32) | Post-Amendment (N=30) |
|--|------------------------|--------------------------|
| Tumor size (cm) | | |
| Median | 4.9 | 4.9 |
| Range | 0.4-14.0 | 0.4-14.0 |
| Histologic grade | | |
| Well differentiated | 2 (6%) | 2 (7%) |
| Moderately differentiated | 14 (44%) | 13 (43%) |
| Poorly differentiated | 7 (22%) | 6 (20%) |
| GX | 9 (28%) | 9 (30%) |
| Perineural invasion | | |
| No | 10 (31%) | 10 (33%) |
| Yes | 22 (69%) | 20 (67%) |
| Lymphovascular invasion | | |
| No | 20 (62%) | 19 (63%) |
| Yes | 12 (38%) | 11 (37%) |
| Margin status | | |
| Negative | 31 (97%) | 29 (97%) |
| Positive | 1 (3%) | 1 (3%) |
| Nodal involvement | | |
| No | 20 (63%) | 20 (67%) |
| Yes | 12 (37%) | 10 (33%) |
| Nodes assessed | | |
| Median | 17 | 16 |
| Range | 4-33 | 4-33 |
| Nodes positive (among patients with nodal involvement) | | |
| Median | 2 | 2 |
| Range | 1-4 | 1-4 |
| Nodes positive, stratified by radiation type | | |
| Long-course | 3 (9%) | 3 (10%) |
| Short-course | 9 (28%) | 7 (23%) |