

Supplementary Online Content

Williamson TJ, Jorge-Miller A, McCannel TA, Beran TM, Stanton AL.
Sociodemographic, medical, and psychosocial factors associated with supportive care needs in adults diagnosed with uveal melanoma. Published online February 22, 2018.
JAMA Ophthalmol. doi:10.1001/jamaophthalmol.2018.0019

eTable. Ten most frequently endorsed unmet supportive care needs

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. *Ten most frequently endorsed unmet supportive care needs*

Rank Order	Item	One Week Post-Diagnosis (%)	Three Months Post-Diagnosis (%)	Domain
1	To be fully informed about cancer remission	87	70	HI
2	To be fully informed about things you can do to help yourself get well	87	66	HI
3	To be fully informed about your test results as soon as possible	86	65	HI
4	Dealing with fears about the cancer spreading or returning	85	61	PSY
5	To be fully informed about the possible effects of the cancer on the length of your life	84	65	HI
6	To be fully informed about the odds of treatment success	83	66	HI
7	To be fully informed about all of the benefits and side effects of treatment or surgery before you agree to have it	80	63	HI
8	To be given a full explanation for every test and treatment procedure you go through	79	61	HI
9	Coping with fears about further physical disability or deterioration	73	48	PSY
10	Coping with fears about the pain and suffering you might experience	72	41	PSY

Note: Unmet needs were analyzed dichotomously as “no need” and “at least 1 unmet need” to evaluate the proportion of unmet needs overall and within each domain. Responses of “not applicable” or “already met” were grouped as “no unmet need” (score = 0) and the low, moderate, and high need responses were grouped as “at least 1 unmet need” (score = 1).

HI = Health Information. PSY = Psychological.