

Supplementary Online Content

Jensen SKG, Dickie EW, Schwartz DH, et al. Effect of early adversity and childhood internalizing symptoms on brain structure in young men. *JAMA Pediatr*. Published online August 17, 2015. doi:10.1001/jamapediatrics.2015.1486.

eAppendix 1. Supplementary Information About Measures

eAppendix 2. Supplementary Information About the Activation Likelihood Estimation (ALE) Definition of Regions of Interest

eAppendix 3. Supplementary Information About the Extraction of Measures of Cortical Gray Matter (GM) Volume, Cortical Thickness and Surface Area in FreeSurfer

eResults. Results From the Univariate and Multivariate Models

eTable 1. Information About the Participant Samples of the Studies Included in the ALE

eTable 2. Results From the Univariate Models

eTable 3. Results From the Final Multivariate Path Model

eFigure. Regions of Interest (ROIs) Identified by the ALE

eReferences.

This supplementary material has been provided by the authors to give readers additional information about their work.

eAppendix 1. Supplementary Information About Measures

Measure of stressful life events

Listed below are the 37 stressful life events included in the assessment of early adversity in the current study. Questionnaires were completed by the mother and collected at six time points after the birth of the child. The questionnaire when the child was 8 months old referred to events since the child was born. The questionnaires collected when the child was 21, 33, 47, 61, and 73 months old referred back to events that had happened within the last year.

Prenatal and adolescent adversity measures were used as control variables. The prenatal questionnaire at 18 weeks gestation referred to events since the mother became pregnant, and did not include questions 33 and 34. Adolescent adversity was measured on a single questionnaire at age 16 years, which referred to experiences while the participant was between 12 and 16 years old.

List of adversities assessed:

1. Mother's partner died
2. A sibling died
3. A family friend died
4. A sibling was ill
5. Mother's partner was ill
6. Mother's friend was ill
7. Mother admitted to hospital
8. Mother in trouble with the law
9. Mother divorced from her partner
10. Mother found out that the child's father did not want the child
11. Mother was very ill
12. Mother's partner lost job
13. Mother's partner had problems with work
14. Mother had problems with work
15. Mother lost her job
16. Mother's partner left her
17. Mother's partner was in trouble with the law
18. Mother was separated from her partner
19. Mother's income was reduced
20. Mother had a big argument with her partner
21. Mother had a big argument with family or close friends
22. The child moved to a new house
23. Mother's partner was cruel to her
24. The family became homeless
25. Mother had financial problems
26. Mother's partner was physically cruel to the child or the child's sibling
27. Mother attempted suicide
28. Mother convicted of an offence
29. Mother started a new job
30. Mother had an abortion
31. Mother's partner was emotionally cruel to the mother
32. Mother's partner was emotionally cruel to the child or the child's sibling
33. Mother was physically cruel to the child or the child's sibling
34. Mother was emotionally cruel to the child or the child's sibling
35. Mother had an accident
36. Mother was under great stress due to an academic examination
37. The family's house was burgled

Measure of internalizing symptom

Pre- and early pubertal levels of internalizing symptoms (depressive/anxiety symptoms) were assessed using the Development and Wellbeing Assessment [DAWBA]. Measures were collected via maternal reports when the boys were 7, 10, and 13 years old. Questions followed the diagnostic criteria in the Diagnostic and Statistical Manual of Mental Disorders [DSM-IV] and the International Classification of Diseases [ICD-10]. We used ordered DAWBA categories that indicate the probability that a child will develop an internalizing disorder on a six-level scale ranging from very unlikely [0.1% of these children have an internalizing disorder] to probable [70% of these children have an internalizing disorder].

eAppendix 2. Supplementary Information About the Activation Likelihood Estimation (ALE) Definition of Regions of Interest

Activation Likelihood Estimation (ALE) is a meta-analytic technique used in the present study as a quantitative way of identifying brain regions that differ in glucose metabolism between patients with depression and healthy controls in a consistent way across multiple studies. The ALE technique was developed originally for meta-analyses of task-related functional neuroimaging (hence “activation” likelihood estimation) but can be applied in the same manner for results obtained with other imaging modalities. The ALE technique assesses the overlap of regions where differences in, for instance, metabolism, have been identified in different studies, by modeling each finding as a probability distribution centered on the reported coordinates.¹ We used results from previously published ¹⁸F-fluorodeoxyglucose Positron Emission Tomography (¹⁸FDG-PET) studies to create a quantitative map of brain regions where lower resting state glucose metabolism has been detected in depressed patients compared with healthy control participants. We used a location-based statistical approach rather than an effect-size based meta-analysis, given that we were more concerned with location than with magnitude of effects. Moreover, the use of coordinates for the ALE-based regions, rather than anatomical labels, minimizes the problem of mislabeling regions due to differences in anatomical referencing between studies.² A similar approach was taken by Sacher et al.,³ yet this previous ALE included just 4 studies, whereas we identified 13 studies (see below for details of the literature search).

Literature search

To conduct the ALE we used existing literature that has identified regions with lower glucose metabolism in depressed patients relative to controls. Such studies were identified via a literature search on PubMed in January 2013. The search combined two statements (#1: depression [mesh] OR depression [ti] OR depression [ab]) and (#2: FDG-PET OR fluorodeoxyglucose). “Mesh” refers to the medical subject headings, “ti” refers to title and “ab” refers to abstracts containing the keywords entered. No time span was specified for date of publication. This search resulted in 842 initial papers. These papers were then examined to ensure the included studies met the following criteria: 1) the study used the ¹⁸F-FDG-PET methodology; 2) the patient population met DSM-IV based diagnostic criteria for depression (bipolar or major depression); 3) the study included control participants; 4) glucose metabolism was measured at rest; 5) the study was not limited to a few preselected ROIs. We did not exclude patients based on medication as this would have seriously limited the number of studies and because the inclusion of medicated patients is commonly accepted in neuroimaging research.⁴ Furthermore, a recent PET study found that lower metabolism observed in cortical regions was indeed more related to disease or mood states than to medication effects.⁵

Fourteen studies met the inclusion criteria for the current ALE⁶⁻¹⁸. Three of these studies^{11,12,18} appeared to have run the relevant analyses, but did not provide the coordinates needed to run the ALE. These authors were contacted and two provided the necessary data upon request.^{12,18} This resulted in inclusion of 13 studies and a sample size of 271 depressive patients and 193 controls in the ALE. eTable 1 summarizes information about the participants of the 13 studies included in the ALE dataset.^{6-10,-12-18}

Subdivision of regions obtained from the ALE analysis

The ALE analysis resulted in a probability map consisting 17 clusters. Large clusters resulting from overlapping peaks/local maxima were further subdivided into smaller clusters with reference to the Automatic Anatomical Labeling atlas¹⁹ using the PickAtlas software²⁰ in SPM8 (Wellcome Department of Imaging Neuroscience, London, UK) implemented in MATLAB 7 (Mathworks Inc., Sherborn, MA). We also split medial ROIs that projected across both hemispheres, into one ROI per hemisphere as FreeSurfer analyses each hemisphere separately. These divisions resulted in 36 clusters/ROIs of which 30 were classified as cortical and 6 were classified as subcortical. This paper focused only on the 30 cortical ROIs, which are shown in the eFigure.

eAppendix 3. Supplementary Information About the Extraction of Measures of Cortical Gray Matter (GM) Volume, Cortical Thickness and Surface Area in FreeSurfer

For every MR image, FreeSurfer segments the cerebral cortex, the white matter, and other subcortical structures, and computes meshes with $\approx 160,000$ triangles that recover the geometry and topology of the pial surface and the gray/white interface of the left and right hemispheres. Regional measures of GM volume, thickness and surface area were obtained by projecting each of the 30 ROIs onto the cortical surface generated using a series of FreeSurfer functions. First, the 30 cortical ROIs were projected from three-dimensional volumes onto the cortical surface of the average brain using *mri_vol2surf*. The 30 surface ROIs were bound into “annotation” files for the left and right hemisphere and then projected from the average template into the participant’s native space using *mri_surf2surf*. Cortical volume, surface area and thickness were then calculated from all ROIs for all participants using *mris_anatomical_stats*. Total brain volume is the total volume of all gray and white matter. Local cortical thickness is measured as a distance between the position of homologous vertices in the pial and gray/white surfaces. A correspondence between the cortical surfaces across participants is established using a nonlinear alignment of the principal sulci in each participant's brain with an average brain.²¹

eResults. Results From the Univariate and Multivariate Models

eTable 2 shows the effects of early adversity and internalizing symptoms on cortical GM volume from the initial univariate models that were part of the model building strategy. Detailed results from the final multivariate path model, which included only the three regions that remained significant after the FDR correction, are found in eTable 3.

eTable 1. Information About the Participant Samples of the Studies Included in the ALE

Authors	N Patients	N Controls	Diagnosis	Age (Mean±SD)		Gender ratio (Male/Female)		Pharmacotherapy?
				Patients	Controls	Patients	Controls	
Broody et al., 2001	24	16	MDD	38.9±11.4	35.6±18.3	13/11	8/8	No
Brooks et al., 2009a	16	11	Bipolar	58.7±7.5	58.3±5.2	14/2	10/1	Yes
Brooks et al., 2009b	15	19	Bipolar	36.1±10.4	34.0±13.3	7/8	12/7	No
Brooks et al., 2010	8	8	Bipolar	52.8±8.5	55.9 ^a	7/1	7/1	Yes
Drevets et al., 1997	17	12	MDD	35±9.4	34±8.2	10/17	11/21	Yes
Hosokawa et al., 2009	47	20	MDD / Bipolar	48.2±16.6	47.6±15.1	0/47	0/20	Mixed
Kegeles et al, 2003	19	10	MDD / Bipolar	36.0±11.0	39.0±19.0	6/13	4/6	Yes
Kennedy et al, 2001	13	24	MDD	36.76±9.37	31.7±6.7	10/0	24/0	No
Martinot et al., 2011	31	39	MDD	47.7±7.3	45.2±11.8	11/20	14/25	Mixed
Mayberg et al., 2001	18	13	MDD	45.0±12.0	38.0±11.0	15/3	12/3	Yes
Yuuki et al., 2005	7	10	MDD	58.0±9.4	60.8±7.9	4/3	5/5	Yes
Suwa et al., 2012	16	11	MDD	32.8±7.9	38.9±10.9	10/6	7/4	Yes
Zhang et al., 2011	40	20	Bipolar	43.4±2.7	N/A	16/24	N/A	Yes
Total	271	193						

N = population size; MDD = Major Depressive Disorder; Bipolar = Bipolar Disorder; SD = Standard Deviation.
 Notes: ^a: SD was not available

eTable 2. Results From the Univariate Models

Effects of early adversity and internalizing symptoms on cortical GM volume. Significant associations ($p < 0.05$) are shown in bold. Control variables are not included in the table.

Risk factor	ROI number	β	SE	<i>P</i> value
L frontopolar cortex (lateral)	1			
Early adversity		-0.05	0.06	.40
Internalizing symptoms		-0.12	0.06	.03
R mid-dorsolateral frontal cortex	2			
Early adversity		0.01	0.05	.95
Internalizing symptoms		-0.05	0.05	.23
L frontopolar cortex (lateral)	3			
Early adversity		-0.08	0.06	.20
Internalizing symptoms		0.04	0.06	.57
R ventromedial prefrontal cortex	4			
Early adversity		-0.04	0.06	.51
Internalizing symptoms		-0.03	0.05	.57
R frontal medial cortex	5			
Early adversity		-0.05	0.07	.45
Internalizing symptoms		-0.07	0.06	.23
L mid-dorsolateral frontal cortex	6			
Early adversity		-0.01	0.06	.85
Internalizing symptoms		-0.03	0.05	.61
L superior frontal sulcus	7			
Early adversity		-0.04	0.06	.53
Internalizing symptoms		-0.08	0.06	.14
R superior frontal gyrus	8			
Early adversity		0.04	0.07	.58
Internalizing symptoms		-0.20	0.07	<.01
R frontal precentral sulcus	9			
Early adversity		0.02	0.07	.81
Internalizing symptoms		-0.03	0.06	.61
R temporal pole	10			
Early adversity		-0.05	0.07	.51
Internalizing symptoms		0.09	0.07	.21
L temporal pole	11			
Early adversity		-0.10	0.08	.17
Internalizing symptoms		0.06	0.07	.41
R superior temporal gyrus	12			
Early adversity		-0.07	0.07	.30
Internalizing symptoms		0.01	0.06	.96
L superior temporal sulcus (anterior)	13			
Early adversity		-0.13	0.07	.06
Internalizing symptoms		0.11	0.055	0.050
R superior temporal sulcus (posterior)	14			
Early adversity		-0.12	0.07	.08
Internalizing symptoms		0.01	0.06	.88
L superior temporal sulcus (posterior)	15			
Early adversity		-0.05	0.07	.51
Internalizing symptoms		0.03	0.06	.59
R supramarginal/angular gyrus	16			
Early adversity		0.16	0.07	.03

Internalizing symptoms		-0.07	0.06	.29
L supramarginal/angular gyrus	17			
Early adversity		0.11	0.08	.17
Internalizing symptoms		-0.12	0.056	0.0
R fusiform gyrus	18			
Early adversity		-0.08	0.07	.30
Internalizing symptoms		-.075	0.061	.22
L fusiform gyrus	19			
Early adversity		-0.14	0.07	.05
Internalizing symptoms		0.020	.060	.736
R lingual gyrus	20			
Early adversity		0.03	0.07	.67
Internalizing symptoms		0.11	0.06	.09
L precuneus	21			
Early adversity		0.02	0.07	.75
Internalizing symptoms		0.07	0.06	.25
R precuneus	22			
Early adversity		0.17	0.07	.01
Internalizing symptoms		-0.05	0.07	.44
L parieto-occipital sulcus	23			
Early adversity		-0.03	0.08	.68
Internalizing symptoms		0.04	0.08	.64
R anterior cingulate cortex (rostral)	24			
Early adversity		-0.05	0.06	.39
Internalizing symptoms		-0.01	0.05	.97
L anterior cingulate cortex (rostral)	25			
Early adversity		-0.06	0.07	.33
Internalizing symptoms		-0.10	0.05	.06
L insula (anterior)	26			
Early adversity		0.05	0.06	.40
Internalizing symptoms		-.06	0.05	.22
R insula (anterior)	27			
Early adversity		0.07	0.07	.33
Internalizing symptoms		-0.04	0.06	.48
L anterior cingulate cortex (caudal)	28			
Early adversity		0.13	0.07	.07
Internalizing symptoms		-.08	0.06	.18
R anterior cingulate cortex (caudal)	29			
Early adversity		-0.17	.07	.01
Internalizing symptoms		0.08	0.07	.21
R parahippocampal gyrus	30			
Early adversity		-0.02	0.06	.77
Internalizing symptoms		0.05	0.07	.45

Vol. = volume, B= standardized path coefficients, SE = Standard Error, *P* = probability value.

eTable 3. Results from the final multivariate path model


Effects of early adversity and internalizing symptoms on cortical GM volume. Significant associations of interest are shown in bold, control variables are shown in italics.

Risk factor	Mean volume [voxels]	SD volume	β	SE	P value
Total brain volume	717985	52360			
2. R superior frontal gyrus	805	170			
Early Adversity			0.05	0.072	.52
Internalizing symptoms			-0.20	0.067	<.01
<i>Total brain volume</i>			<i>0.34</i>	<i>0.040</i>	<i><.01</i>
<i>Adolescent adversity (12-16y)</i>			<i>0.04</i>	<i>0.048</i>	<i>.39</i>
<i>Prenatal family stress</i>			<i>0.06</i>	<i>.057</i>	<i>.30</i>
<i>Duration breast feeding</i>			<i>-0.04</i>	<i>0.045</i>	<i>.35</i>
5. R precuneus	992	196			
Early Adversity			0.18	0.07	<.01
Internalizing symptoms			-0.05	0.07	.41
<i>Total brain volume</i>			<i>0.38</i>	<i>0.039</i>	<i><0.001</i>
<i>Adolescent adversity (12-16y)</i>			<i>0.02</i>	<i>0.05</i>	<i>.70</i>
<i>Prenatal family stress</i>			<i>-0.11</i>	<i>0.06</i>	<i>.06</i>
<i>Duration breast feeding</i>			<i>0.03</i>	<i>0.05</i>	<i>.46</i>
6. R Anterior cingulate cortex	642	180			
Early Adversity			-0.18	0.07	.01
Internalizing symptoms			0.08	.07	.31
<i>Total brain volume</i>			<i>0.27</i>	<i>0.04</i>	<i><.01</i>
<i>Adolescent adversity (12-16y)</i>			<i>0.004</i>	<i>0.05</i>	<i>.94</i>
<i>Prenatal family stress</i>			<i>0.04</i>	<i>0.06</i>	<i>.50</i>
<i>Duration breast feeding</i>			<i>-0.08</i>	<i>0.05</i>	<i>.08</i>

SD = Standard Deviation, B= standardized path coefficients, SE = Standard Error, P = probability value.

eFigure. Regions of Interest (ROIs) Identified by the ALE

Image showing the 30 cortical regions of interest (ROIs) identified by the ALE analysis and separated into smaller regions using PickAtlas. The ROIs are shown on transverse slices ranging from z=-35 to z=65 (Montreal Neurological Institute coordinates). Yellow numbers refer to the ROI numbers presented in Table 1 in the article and eTable 2.



eReferences

1. Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT (2009): Coordinate-based activation likelihood estimation meta-analysis of neuroimaging data: A random-effects approach based on empirical estimates of spatial uncertainty. *Hum Brain Mapp.* 30:2907-2926.
2. Fusar-Poli P, Placentino A, Carletti F, Landi P, Allen P, Surguladze S, et al. (2009): Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *J Psychiatry Neurosci.* 34:418-432.
3. Sacher J, Neumann J, Fünfstück T, Soliman A, Villringer A, Schroeter ML (2012): Mapping the depressed brain: a meta-analysis of structural and functional alterations in major depressive disorder. *J Affect Disord.* 140:142-148.
4. Phillips M, Travis M, Fagiolini A, Kupfer D (2008): Medication effects in neuroimaging studies of bipolar disorder. *Am J Psychiatry.* 165:313-320.
5. Gonul AS, Coburn K, Kula M (2009): Cerebral blood flow, metabolic, receptor, and transporter changes in bipolar disorder: the role of PET and SPECT studies. *Int Rev Psychiatry.* 21:323-335.
6. Brody, A. L., Saxena, S., Mandelkern, M. A., Fairbanks, L. A., Ho, M. L., & Baxter, L. R. (2001). Brain metabolic changes associated with symptom factor improvement in major depressive disorder. *Biol psychiatry.* 50(3), 171-178.
7. Brooks JO, Hoblyn JC, Ketter TA (2010): Metabolic evidence of corticolimbic dysregulation in bipolar mania. *Psychiatry Res.* 181:136-140.
8. Brooks JO, Hoblyn JC, Woodard SA, Rosen AC, Ketter TA (2009): Corticolimbic metabolic dysregulation in euthymic older adults with bipolar disorder. *J Psychiatr Res.* 43:497-502.
9. Brooks JO, Wang PW, Bonner JC, Rosen AC, Hoblyn JC, Hill SJ, et al. (2009): Decreased prefrontal, anterior cingulate, insula, and ventral striatal metabolism in medication-free depressed outpatients with bipolar disorder. *J Psychiatric Res.* 43:181-188.
10. Drevets WC, Price JL, Simpson JR, Todd RD, Reich T, Vannier M, et al. (1997): Subgenual prefrontal cortex abnormalities in mood disorders. *Nature.* 386:824-828.
11. Fujimoto T, Takeuchi K, Matsumoto T, Fujita S, Honda K, Higashi Y, et al. (2008): Metabolic changes in the brain of patients with late-onset major depression. *Psychiatry Res.* 164:48-57.
12. Hosokawa T, Momose T, Kasai K (2009): Brain glucose metabolism difference between bipolar and unipolar mood disorders in depressed and euthymic states. *Prog Neuropsychopharmacol Biol Psychiatry.* 33:243-250.
13. Kegeles LS, Malone KM, Slifstein M, Ellis SP, Xanthopoulos E, Keilp JG, et al. (2003): Response of cortical metabolic deficits to serotonergic challenge in familial mood disorders. *Am J Psychiatry.* 160:76-82.
14. Kennedy SH, Evans KR, Krüger S, Mayberg HS, Meyer JH, McCann S, et al. (2001): Changes in regional brain glucose metabolism measured with positron emission tomography after paroxetine treatment of major depression. *Am J Psychiatry.* 158:899-905.
15. Mayberg HS, Brannan SK, Mahurin RK, Jerabek PA, Brickman JS, Tekell JL, et al. (1997): Cingulate function in depression: a potential predictor of treatment response. *Neuroreport.* 8:1057-1061.
16. Paillere Martinot ML, Martinot JL, Ringuenet D, Galinowski A, Gallarda T, Bellivier F, et al. (2011): Baseline brain metabolism in resistant depression and response to transcranial magnetic stimulation. *Neuropsychopharmacology.* 36:2710-2719.
17. Yuuki N, Ida I, Oshima A, Kumano H, Takahashi K, Fukuda M, et al. (2005): HPA axis normalization, estimated by DEX/CRH test, but less alteration on cerebral glucose metabolism in depressed patients receiving ECT after medication treatment failures. *Acta Psychiatr Scand.* 112:257-265.
18. Zhang L, Li CT, Su TP, Hu XZ, Lanius RA, Webster MJ, et al. (2011): P11 expression and PET in bipolar disorders. *J Psychiatr Res* 45:1426-1431.

19. Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, et al. (2002): Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*. 15:273-289.
20. Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH (2003): An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *NeuroImage*. 19:1233-1239.
21. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. (2006): An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*. 31:968-980.