

## Supplementary Online Content

Keyes KM, Platt J, Kaufman AS, McLaughlin KA. Association of fluid intelligence and psychiatric disorders in a population-representative sample of US adolescents. *JAMA Psychiatry*. Published online December 28, 2016. doi:10.1001/jamapsychiatry.2016.3723

**eTable 1.** Sample sizes for each disorder group used in the analysis, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months

**eTable 2.** Unadjusted mean differences in IQ by psychiatric disorder among US adolescents (N=10,073)

**eTable 3.** Effect size estimates of the association between fluid intelligence and psychiatric disorders in a population representation sample of adolescents (N=10,073)

**eTable 4.** Variation in fluid intelligence by current disorder status, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months

**eTable 5.** Variation in fluid intelligence as a function of lifetime psychiatric disorders (N=10,073)

**eTable 6.** Proportion of adolescents with fluid intelligence in low, middle, and high range, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months

**eTable 7.** Unadjusted mean differences in IQ by psychiatric disorder among US adolescents by disorder age of onset

**eTable 8.** Proportion of adolescents with fluid intelligence in low, middle, and high range as a function of psychiatric disorder age of onset (unadjusted) (N=10,073)

**eTable 9.** Adjusted IQ differences as a function of psychiatric disorder age of onset in a population representation sample of adolescents (N=10,073)

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Sample sizes for each disorder group used in the analysis, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months								
	<i>Past 12 month and prior to past 12 months, all ages</i>				<i>Past 12-month but not prior to past 12 months, all ages</i>			
	Total	Low (%)	Middle (%)	High (%)	Total	Low (%)	Middle (%)	High (%)
<i>I. Fear disorders</i>								
Specific phobia	1600	360	1042	198	21	5	16	--
Agoraphobia	217	55	126	36	5	3	2	--
Social phobia	1273	256	848	169	--	--	--	--
Panic disorder	191	35	136	20	--	--	--	--
No lifetime fear disorder	7164	1249	4824	1091	7164	1249	4824	1091
<i>II. Distress disorders</i>								
Separation anxiety disorder	162	42	107	13	12	4	7	1
Post-traumatic stress disorder	288	50	194	44	11	3	7	1
Major depressive episode/dysthymia	949	182	638	129	47	12	32	3
Generalized anxiety disorder	176	41	111	24	11	3	7	1
No lifetime distress disorder	4473	818	3036	619	4473	818	3036	619
<i>III. Behavior disorders</i>								
ADHD	247	70	149	28	2	1	1	--
Oppositional defiant disorder	488	113	328	47	49	10	32	7
Conduct disorder	333	90	216	27	91	17	67	7
Eating disorders	311	76	195	40	--	--	--	--
No lifetime behavior disorder	8103	1401	5483	1219	8103	1401	5483	1219
<i>IV. Substance disorders</i>								
Alcohol abuse	504	110	344	50	42	11	26	5
Drug abuse	548	107	376	65	100	15	76	9
No lifetime substance disorder	8912	1620	5951	1341	8912	1620	5951	1341
<i>V. Other disorders</i>								
Bipolar disorder	113	32	69	12	5	1	3	1
No lifetime bipolar disorder	8831	1591	5905	1335	8831	1591	5905	1335

eTable 2. Unadjusted mean differences in IQ by psychiatric disorder among US adolescents (N=10,073)						
	<i>Past 12 month disorders, all ages</i>			<i>Prior to past 12-month but not current disorder, all ages</i>		
	Mean IQ	SD	p value <sup>+</sup>	Mean IQ	SD	p value <sup>+</sup>
<i>I. Fear disorders</i>						
Specific phobia	96.6	16.4	0.004	99.9	14.3	0.91
Agoraphobia	97.3	14.8	0.002	97.8	15.4	0.47
Social phobia	98.4	14.6	0.004	98	15.4	0.47
Panic disorder	97.9	13.9	0.10	96.8	14.3	0.47
No fear disorder	100.1	14.5	--	100.1	14.5	--
<i>II. Distress disorders</i>						
Separation anxiety disorder	95	14.5	0.30	97	14.6	0.002
Post-traumatic stress disorder**	100	14.3	0.30	96.7	14.6	0.17
Major depressive episode/dysthymia	99.5	14.6	0.92	100	14	0.55
Generalized anxiety disorder	98.1	16.1	0.92	97.4	13.3	0.17
No distress disorder	98.1	14.4	--	99.8	14.6	--
<i>III. Behavior disorders</i>						
ADHD	96.1	16.1	0.002	96.2	14	0.004
Oppositional defiant disorder	96.6	14.9	<.001	98.9	15	0.14
Conduct disorder	95.5	14.7	<.001	96.2	14.6	0.004
Eating disorders	97	15.1	0.002	98.2	15	0.14
No behavior disorder	100	14.5	--	100.1	14.5	--
<i>IV. Substance disorders</i>						
Alcohol abuse	97.4	14.8	0.004	97.9	14.8	0.11
Drug abuse	98.3	14.3	0.03	98.3	13.2	0.11
No substance disorder	99.8	14.6	--	99.8	14.6	--
<i>V. Other disorders</i>						
Bipolar disorder	94.8	16.2	0.001	97.1	16.3	0.41
No bipolar disorder	99.6	14.6	--	99.7	14.6	--
<i>VI. Total number of disorders</i>						
Exactly one disorder	98.7	13.6	0.40	100.5	14.5	0.10
Exactly two disorders	98.9	13.8	0.40	97.7	13.7	0.10
Three or more disorders	97.6	14.9	0.02	96.6	14.3	0.10
No disorder	99.7	14.6	--	99.6	14.6	--
* Scores were first normed in the sample by six-month age groups for mean of 100 and standard deviation of 15.						
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample)						
+ P-values are for the comparison between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear disorder. All p-values are false discovery rate adjusted.						

eTable 3. Effect size estimates of the association between fluid intelligence and psychiatric disorders in a population representation sample of adolescents (N=10,073)*					
	<i>Past 12 month disorders, all ages</i>		<i>Prior to past 12-month but not current disorder, all ages</i>		
	d	CI	d	CI	
<i>I. Fear disorders</i>					
Specific phobia	0.12	0.07, 0.18	0.00	-0.11, 0.11	
Agoraphobia	0.02	-0.12, 0.16	0.06	-0.17, 0.29	
Social phobia	0.03	-0.03, 0.09	0.11	-0.05, 0.27	
Panic disorder	0.05	-0.09, 0.19	0.15	-0.15, 0.45	
<i>II. Distress disorders</i>					
Separation anxiety disorder	0.14	-0.02, 0.29	0.11	-0.19, 0.41	
Post-traumatic stress disorder**	-0.06	-0.18, 0.06	0.14	-0.06, 0.34	
Major depressive episode/dysthymia	-0.08	-0.15, -0.01	-0.06	-0.16, 0.04	
Generalized anxiety disorder	0.07	-0.08, 0.23	0.16	-0.02, 0.34	
<i>III. Behavior disorders</i>					
ADHD	0.17	0.05, 0.3	0.14	-0.01, 0.29	
Oppositional defiant disorder	0.12	0.03, 0.21	0.03	-0.06, 0.12	
Conduct disorder	0.14	0.03, 0.25	0.10	-0.03, 0.23	
Eating disorders	0.08	-0.03, 0.19	0.05	-0.08, 0.18	
<i>IV. Substance disorders</i>					
Alcohol abuse	0.18	0.09, 0.27	0.10	-0.05, 0.25	
Drug abuse	0.10	0.02, 0.19	0.09	-0.02, 0.2	
<i>V. Other disorders</i>					
Bipolar disorder	0.31	0.12, 0.49	0.06	-0.36, 0.48	
<i>VI. Total number of disorders</i>					
Exactly one disorder	0.04	-0.07, 0.16	-0.06	-0.17, 0.05	
Exactly two disorders	0.04	-0.09, 0.18	0.09	-0.04, 0.22	
Three or more disorders	0.07	-0.02, 0.16	0.20	0.11, 0.29	
*Adjusted for parental education, parental income, poverty ratio, race/ethnicity, age, and nativity, number of siblings, birth order, and non-focal disorders					
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample).					

eTable 4. Variation in fluid intelligence* by current disorder status, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months								
	<i>Past 12 month and Lifetime disorders, all ages</i>				<i>Past 12 month, but not Lifetime disorders, all ages</i>			
	Mean IQ	SE	Beta	p value+	Mean IQ	SE	Beta	p value+
<i>I. Fear disorders</i>								
Specific phobia	97.9	0.39	-1.3	0.005	96.7	3.11	-2.49	0.42
Agoraphobia	98.8	0.98	-0.45	0.65	87.5	6.27	-11.8	0.12
Social phobia	98.6	0.43	-0.51	0.50	--	--	--	--
Panic disorder	98.4	1.04	-0.9	0.52	--	--	--	--
No fear disorder	99.1	0.22	--	--	99.1	0.22	--	--
<i>II. Distress disorders</i>								
Separation anxiety disorder	96.8	1.13	-1.89	0.20	98.3	4.19	-0.41	0.97
Post-traumatic stress disorder**	99.7	0.88	0.94	0.32	98.6	4.38	-0.19	0.97
Major depressive episode/dysthymia	100	0.5	1.32	0.04	96.8	2.04	-1.91	0.70
Generalized anxiety disorder	97.6	1.11	-1.11	0.32	93.2	4.16	-5.53	0.67
No distress disorder	98.8	0.25	--	--	98.8	0.25	--	--
<i>III. Behavior disorders</i>								
ADHD	96.3	0.91	-2.91	0.006	93.6	9.8	-5.63	0.90
Oppositional defiant disorder	97.3	0.66	-1.81	0.01	99.2	2	0.1	0.96
Conduct disorder	97.1	0.82	-1.96	0.02	98.3	1.46	-0.77	0.90
Eating disorders	97.9	0.82	-1.28	0.12	--	--	--	--
No behavior disorder	99.1	0.21	--	--	99.1	0.21	--	--
<i>IV. Substance disorders</i>								
Alcohol abuse	96.5	0.67	-2.6	<.001	97.6	2.16	-1.47	0.93
Drug abuse	97.6	0.64	-1.45	0.02	98.9	1.41	-0.13	0.93
No substance disorder	99.1	0.21	--	--	99.1	0.21	--	--
<i>V. Other disorders</i>								
Bipolar disorder	94.3	1.69	-4.88	0.004	106.6	6.3	7.4	0.24
No bipolar disorder	99.2	0.21	--	--	99.2	0.21	--	--
<i>VI. Total number of disorders</i>								
Exactly one disorder	98.1	0.34	-1.32	<.001	97.8	0.77	-1.03	0.05
Exactly two disorders	98.1	0.48	-1.37	0.006	97.6	2.98	-1.29	0.66
Three or more disorders	97	0.49	-2.48	<.001	90.5	9.86	-8.38	0.59
No disorder	99.6	0.25	--	--	99.6	0.25	--	--
* Scores were first normed in the sample by six-month age groups for mean of 100 and standard deviation of 15. Predicted means were estimated from linear regression models controlling for parental education; race/ethnicity, age, nativity (US born versus not), number of siblings, birth order, and non-focal disorder groups								
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample)								
+ P-values are for the comparison between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear								

disorder. All p-values are false discovery rate adjusted.

eTable 5. Variation in fluid intelligence* as a function of lifetime psychiatric disorders (N=10,073)					
	Mean IQ	SE	p value <sup>+</sup>	d	CI
<i>I. Fear disorders</i>					
Specific phobia	97.7	0.35	0.004	0.1	0.05, 0.14
Agoraphobia	98.2	0.83	0.44	0.06	-0.06, 0.17
Social phobia	98.1	0.4	0.07	0.07	0.01, 0.12
Panic disorder	97.3	0.93	0.14	0.13	0, 0.26
No fear disorder	99.1	0.22	--	--	
<i>II. Distress disorders</i>					
Separation anxiety disorder	97	0.53	0.004	0.12	0.03, 0.22
Post-traumatic stress disorder**	98.2	0.75	0.58	0.04	-0.08, 0.16
Major depressive episode/dysthymia	99.1	0.41	0.52	-0.02	-0.1, 0.06
Generalized anxiety disorder	96.7	0.83	0.02	0.14	0.01, 0.27
No distress disorder	99	0.21	--	--	
<i>III. Behavior disorders</i>					
ADHD	96.1	0.69	<.001	0.2	0.11, 0.3
Oppositional defiant disorder	97.5	0.46	0.003	0.11	0.05, 0.18
Conduct disorder	96.4	0.6	<.001	0.19	0.11, 0.27
Eating disorders	97.6	0.61	0.05	0.11	0.02, 0.19
No behavior disorder	99.2	0.21	--	--	
<i>IV. Substance disorders</i>					
Alcohol abuse	96.2	0.58	<.001	0.2	0.12, 0.27
Drug abuse	97.3	0.51	0.001	0.13	0.06, 0.2
No substance disorder	99.1	0.21	--	--	
<i>V. Other disorders</i>					
Bipolar disorder	95.55	1.22	0.007	0.21	0.04, 0.38
No bipolar disorder	98.9	0.2	--	--	
<i>VI. Total number of disorders</i>					
No disorder	98.8	0.21	--	--	
Exactly one disorder	98.7	0.42	0.17	0.04	-0.08, 0.15
Exactly two disorders	98.7	0.52	0.22	0.04	-0.1, 0.17
Three or more disorders	97.7	0.37	<.001	0.1	0.03, 0.18
* Scores were first normed in the sample by six-month age groups for mean of 100 and standard deviation of 15. Predicted means were estimated from linear regression models controlling for parental education; race/ethnicity, age, nativity (US born versus not), number of siblings, birth order, and non-focal disorder groups					
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample)					
<sup>+</sup> P-values are for the comparison between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear disorder. All p-values are false discovery rate adjusted.					

eTable 6. Proportion of adolescents with fluid intelligence in low, middle, and high range, separating current disorders into those with onset in the past 12 months and those with onset prior to the past 12 months							
	<i>Past 12 month and Lifetime disorders, all ages</i>				<i>Past 12 months but not lifetime, all ages</i>		
<i>Range (N total)</i>	Low (n=1852) (%)	Middle (n=6757) (%)	High (n=1464) (%)	p-value <sup>+</sup>	Middle (n=6757) (%)	High (n=1464) (%)	p-value <sup>+</sup>
<i>I. Fear disorders</i>							
Specific phobia	22.5	65.13	12.38	<.001	76.19	0	0.14
Agoraphobia	25.35	58.06	16.59	0.009	40	0	0.08
Social phobia	20.11	66.61	13.28	0.035	0	0	--
Panic disorder	18.32	71.2	10.47	0.19	0	0	--
Any fear disorder	21.02	66.13	12.85	<.001	69.23	0	0.08
No fear disorder	17.52	67.39	15.09	--	67.07	14.57	--
<i>II. Distress disorders</i>							
Separation anxiety disorder	25.93	66.05	8.02	0.06	58.33	8.33	0.64
Post-traumatic stress disorder**	17.36	67.36	15.28	0.76	63.64	9.09	0.71
Major depressive episode/dysthymia	19.18	67.23	13.59	0.76	68.09	6.38	0.44
Generalized anxiety disorder	23.3	63.07	13.64	0.39	63.64	9.09	0.71
Any distress disorder	20.24	66.48	13.28	0.29	65	7.5	0.20
No distress disorder	18.12	67.17	14.71	--	67.1	14.59	--
<i>III. Behavior disorders</i>							
ADHD	28.34	60.32	11.34	<.001	50	0	0.59
Oppositional defiant disorder	23.16	67.21	9.63	<.001	65.31	14.29	0.85
Conduct disorder	27.03	64.86	8.11	<.001	73.63	7.69	0.58
Eating disorders	24.44	62.7	12.86	0.005	0	0	--
Any behavior disorder	24.64	64.55	10.8	<.001	70.71	10	0.59
No behavior disorder	17.6	67.4	15	--	67.03	14.6	--
<i>IV. Substance disorders</i>							
Alcohol abuse	21.83	68.25	9.92	0.007	61.9	11.9	0.39
Drug abuse	19.53	68.61	11.86	0.12	76	9	0.36
Any substance disorder	20.47	68.59	10.94	0.007	71.85	9.63	0.38
No substance disorder	18.19	66.94	14.87	--	67.02	14.6	--
<i>V. Other disorders</i>							



Bipolar disorder	28.32	61.06	10.62	0.01	60	20	0.94
No bipolar disorder	18.02	66.87	15.12	--	66.87	15.12	--
<i>VI. Total number of disorders</i>							
Exactly one disorder	19.51	67.38	13.11	0.00 1	70.23	8.96	0.02
Exactly two disorders	21.58	64.97	13.45	0.00 1	68.18	9.09	0.70
Three or more disorders	23.28	65.92	10.8	<.00 1	50	0	0.64
Any disorder	20.88	66.47	12.66	<.00 1	70	8.92	0.02
No disorder	16.84	67.46	15.7	--	66.97	14.75	--
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample)							
† P-values are for chi-square comparisons between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear disorder. All p-values are false discovery rate adjusted.							

eTable 7. Unadjusted mean differences in IQ by psychiatric disorder among US adolescents by disorder age of onset									
	Onset age 4-8			Onset age 9-12			Onset age 13-17		
	Mean IQ	SD	p value <sup>+</sup>	Mean IQ	SD	p value <sup>+</sup>	Mean IQ	SD	p value <sup>+</sup>
<i>I. Fear disorders</i>									
Specific phobia	98.1	14.6	<.001	96.3	15.1	<.001	96.9	16	0.19
Agoraphobia	97.6	15.7	0.08	96.6	17	0.02	95.6	15.5	0.06
Social phobia	98.4	14.4	0.01	99.3	15.1	0.15	95.6	13.9	<.001
Panic disorder	99.7	13.3	0.79	96.3	13.6	0.02	97.1	14.9	0.11
No fear disorder	100.1	14.5	--	100.1	14.5	--	100.1	14.5	--
<i>II. Distress disorders</i>									
Separation anxiety disorder	96.9	14.8	<.001	96.8	13.9	0.02	94.6	15.5	0.004
Post-traumatic stress disorder**	99	13.4	0.89	98.8	16.4	0.78	99.5	13.8	0.87
Major depressive episode/dysthymia	99.4	15	0.89	100.5	14	0.13	99.1	14.6	0.87
Generalized anxiety disorder	98.5	14.1	0.89	99.9	13.2	0.78	96	16.4	0.02
No distress disorder	99.2	14.5	--	99.2	14.5	--	99.2	14.5	--
<i>III. Behavior disorders</i>									
ADHD	97.6	14.7	0.01	92.3	16.3	<.001	89	15.5	0.002
Oppositional defiant disorder	98.1	15.3	0.03	98.2	14.6	0.02	97.1	15.4	0.002
Conduct disorder	102.5	13.6	0.03	98.7	14.5	0.07	97.5	14.6	0.003
Eating disorders	95	16.4	0.03	97.4	15.2	0.02	97.9	14.9	0.009
No behavior disorder	100.1	14.5	--	100.1	14.5	--	100.1	14.5	--
<i>IV. Substance disorders</i>									
Alcohol abuse	109.8	11.1	0.18	97.4	14.7	0.40	97.5	14.8	0.002
Drug abuse	84.4	3.8	0.07	98.4	13.7	0.40	98.3	13.9	0.007
No substance disorder	99.8	14.6	--	99.8	14.6	--	99.8	14.6	--
<i>V. Other disorders</i>									
Bipolar disorder	92.7	15.9	0.03	96.3	16.9	0.12	95.1	15.8	0.01
No bipolar disorder	99.7	14.6	--	99.7	14.6	--	99.7	14.6	--
** Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of									

the total sample)		
+ P-values are for the comparison between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear disorder. All p-values are false discovery rate adjusted.		

eTable 8. Proportion of adolescents with fluid intelligence in low, middle, and high range as a function of psychiatric disorder age of onset (unadjusted) (N=10,073)												
	Onset age 4-8				Onset age 9-12				Onset age 13-17			
	Low (%)	Middle (%)	High (%)	p-value <sup>†</sup>	Low (%)	Middle (%)	High (%)	p-value <sup>†</sup>	Low (%)	Middle (%)	High (%)	p-value <sup>†</sup>
<i>I. Fear disorders</i>												
Specific phobia	20.76	66.58	12.65	****	24.13	64.76	11.11	**	25.71	65.71	8.57	*
Agoraphobia	22.4	60	17.6	*	24.79	60.68	14.53	*	33.33	52.08	14.58	**
Social phobia	20.14	67.45	12.41	*	19.69	64.22	16.09	*	23.66	67.41	8.93	**
Panic disorder	19.23	70.51	10.26	*	19.77	69.77	10.47	*	21.13	69.01	9.86	*
Any fear disorder	20.73	66.34	12.82	****	20.73	66.34	12.82	****	20.73	66.34	12.82	****
No fear disorder	17.43	67.34	15.23	--	17.43	67.34	15.23	--	17.43	67.34	15.23	--
<i>II. Distress disorders</i>												
Separation anxiety disorder	23.35	64.88	11.78	***	21.39	68.45	10.16	*	33.7	56.52	9.78	****
Post-traumatic stress disorder <sup>1</sup>	17.76	70.09	12.15	*	21.57	58.82	19.61	*	18.97	67.82	13.22	*
Major depressive episode/dysthymia	18.36	68.12	13.53	*	16.36	69.09	14.55	*	20.31	66.11	13.59	
Generalized anxiety disorder	15.15	71.21	13.64	*	14.29	73.47	12.24	*	31.58	54.14	14.29	****
Any distress disorder	18.46	66.45	15.09	*	18.46	66.45	15.09	*	18.46	66.45	15.09	*
No distress disorder	18.29	67.87	13.84	--	18.29	67.87	13.84	--	18.29	67.87	13.84	--
<i>III. Behavior disorders</i>												
ADHD	23.24	65.44	11.31	**	33.33	58.97	7.69	***	52.17	39.13	8.7	****
Oppositional defiant disorder	20.07	67.07	12.59	*	21.4	66.51	12.09	*	24.84	61.64	13.52	**
Conduct disorder	11.41	70.11	18.48	**	20.05	67.45	12.5	*	20.88	68.01	11.11	*
Eating disorders	35.71	47.62	16.67	***	23.71	65.46	10.82	*	20.89	65.51	13.61	*
Any behavior disorder	22.89	64.67	12.44	****	22.89	64.67	12.44	****	22.89	64.67	12.44	****
No behavior disorder	17.29	67.67	15.04	--	17.29	67.67	15.04	--	17.29	67.67	15.04	--
<i>IV. Substance disorders</i>												
Alcohol abuse	--	75	25	--	21.57	70.59	7.84	*	21.81	68.01	10.18	***
Drug abuse	50	50	--	--	20.99	72.84	6.17	*	19.22	69.41	11.38	**

Any substance disorder	19. 98	69.4 2	10. 59	****	19. 98	69.4 2	10. 59	****	19. 98	69.4 2	10. 59	****
No substance disorder	18. 18	66.7 8	15. 05	--	18. 18	66.7 8	15. 05	--	18. 18	66.7 8	15. 05	--
<i>V. Other disorders</i>												
Bipolar disorder	35	55	10	*	25. 53	63.8 3	10. 64	*	27. 94	63.2 4	8.8 2	*
No bipolar disorder	18. 25	67.1 5	14. 6	--	18. 25	67.1 5	14. 6	--	18. 25	67.1 5	14. 6	--
<sup>1</sup> Among those with a lifetime exposure to a potentially traumatic event (N=6160, 61.2% of the total sample)												
<sup>+</sup> P-values are for chi-square comparisons between each disorder category to a reference group of no disorder in that category. For example, mean IQ among those with specific phobia is compared to those with no fear disorder. All p-values are false discovery rate adjusted. * = p<0.05; ** = p<0.05; *** = p<0.01; **** = p<0.001												

eTable 9. Adjusted IQ differences as a function of psychiatric disorder age of onset in a population representation sample of adolescents (N=10,073)									
	Onset age 4-8			Onset age 9-12			Onset age 13-17		
	Mean IQ	SE	p value <sup>+</sup>	Mean IQ	SE	p value <sup>+</sup>	Mean IQ	SE	p value <sup>+</sup>
<i>I. Fear disorders</i>									
Specific phobia	98.2	0.38	0.56	97.6	0.8	0.63	96.6	2.4	0.80
Agoraphobia	99.1	1.27	0.56	98.6	1.32	0.63	97.9	2.03	0.80
Social phobia	98.3	0.61	0.02	99.3	0.58	0.35	96.4	0.96	0.03
Panic disorder	99.9	1.6	0.56	96.3	1.53	0.63	98.3	1.67	0.75
No fear disorder	99.1	0.22	--	99.1	0.22	--	99.1	0.22	--
<i>II. Distress disorders</i>									
Separation anxiety disorder	97.4	0.66	0.08	98.1	1.04	0.68	95.9	1.46	0.28
Post-traumatic stress disorder**	98.5	1.37	0.70	99.1	1.41	0.68	99.2	1.12	0.96
Major depressive episode/dysthymia	99.8	1.01	0.70	100.2	0.67	0.68	99.1	0.59	0.28
Generalized anxiety disorder	98.8	1.76	0.04	98.6	1.44	0.47	95.5	1.25	0.28
No distress disorder	98.8	0.25	--	98.8	0.25	--	98.8	0.25	--
<i>III. Behavior disorders</i>									
ADHD	97.5	0.8	0.11	94.2	1.6	0.21	92.8	2.9	0.67
Oppositional defiant disorder	97.6	0.84	0.15	98.6	0.69	0.94	97.7	0.81	0.67
Conduct disorder	101.1	1.03	0.40	98.9	0.73	0.15	98.4	0.83	0.67
Eating disorders	95.4	2.14	0.50	97.9	1.02	0.21	98.7	0.82	0.67
No behavior disorder	99.1	0.21	--	99.1	0.21	--	99.1	0.21	--
<i>IV. Substance disorders</i>									
Alcohol abuse	110.7	6.98	<.001	97.1	1.97	0.09	96.6	0.62	0.81
Drug abuse	87.1	6.95	0.01	98.6	1.57	0.13	97.7	0.55	0.81
No substance disorder	99	0.21	--	99	0.21	--	99	0.21	--
<i>V. Other disorders</i>									
Bipolar disorder	94	3.12	0.16	98.4	2.05	0.49	96.4	1.73	0.47
No bipolar disorder	98.9	0.2	--	98.9	0.2	--	98.9	0.2	--