

## Supplementary Online Content


Wilson DG, Harris SK, Peck H, et al. Patterns of care in hospitalized vascular surgery patients at end of life. *JAMA Surg*. Published online November 2, 2016.  
doi:10.1001/jamasurg.2016.3970

**eFigure.** Flow diagram – patient selection

**eTable.** Multivariable logistic regression for comfort care with other medical options available vs controls

This supplementary material has been provided by the authors to give readers additional information about their work.

**eFigure. Flow diagram – patient selection**



**eTable. Multivariable logistic regression for comfort care with other medical options available vs controls**

	OR (95% CI)	p-value
More than 5 days on ventilator or More than 5 days in ICU	3.26 (0.72, 14.70)	0.12
Trach, Dialysis, or CPR	14.20 (3.16, 63.71)	<0.01
Emergency vs. Elective Admission	2.63 (0.56, 12.25)	0.22
Obesity BMI >30	0.62 (0.12, 3.26)	0.57
Family communication at admission	0.85 (0.11, 6.30)	0.87
Time of operation (6:01pm-12am vs. 6am-6pm)	0.45 (0.04, 4.83)	0.51
Time of operation (12am-5:59am vs. 6am-6pm)	1.95 (0.29, 13.22)	0.49
Hosmer DW, Lemeshow S. Applied Logistic Regression. New York: Wiley;2000		